

Utility of Thermotherapy and Meristem Tip for Freeing Sweetpotato from Viral Infection.

Mervat M. M. El Far and A. Ashoub

Agricultural Genetic Engineering Research Institute (AGERI), Agricultural Research Centre (ARC),
9 Gamaa St. 12619, Giza, Egypt

Abstract: Sweetpotato, (*Ipomoea batatas L.*), is vegetatively propagated crop. Due to the following cycles of propagation, viruses are accumulated, which contributes to the decline of sweetpotato yield and quality. Stem cuttings of Abees cultivar were collected from several locations in Egypt and kept under controlled greenhouse conditions. Samples were tested for the presence of sweetpotato feathery mottle virus (SPFMV) infection using dot-ELISA. Infected plants were thermotherapy treated by incubation of plants at 42°C/ day and 39°C/ night for 3 weeks followed by meristem tip culturing and allowed to grow *in vitro*. RT-PCR was carried out to confirm the success of SPFMV elimination. Tissue culture formed plants were tested routinely for successive 2 years using dot-ELISA. 0% infection was reported in the *in vitro* propagated plants. Sweetpotato plants were assisted the private-sector to enhance the final yield in their bio-farming system.

Key words: sweetpotato; virus-free; dot-ELISA; thermotherapy.

INTRODUCTION

Sweetpotato (*Ipomoea batatas L.*) is a dicotyledonous, perennial plant, belongs to the family convolvulacea (Astin D.F., 1987). It ranks as the seventh important staple crop in the world and the fifth in developing countries after rice, wheat, maize and cassava (Loebenstein G., *et al.*, 2003). Sweetpotato considered being the second important root crop after cassava in many tropical countries (FAOSTAT, 2006). Because of the enormous genetic diversity of sweetpotato (Zhang D., *et al.*, 1998), and the accompanying diversity in phenotypic and morphological traits (Woolfe J.A., 1992), the crop has great potential for further development to accommodate specific uses. On the other hand, nutritional studies in Egypt gave the evidence that vitamin A deficiency (VAD) is prevalent among pre-school children and their mothers (Nutrition Institute, 1995). VAD is a main cause of child blindness, and even in its less acute forms, it hinders normal growth and development (WHO nutrition report, 2003). Orange fleshed sweetpotato varieties are rich in B-carotene, the precursor of vitamin A, (CIP, 2001). In Egypt, the sweetpotato variety Abees, a purple skin and orange flesh, has a high productivity per unit/area, good nutritional value and a big demand for exportation. Abees cultivar represents approximately 60% of the total sweetpotato acreage (30.000 acres).

Despite of the advantages that the cultivation of sweetpotato offers, production is greatly constrained particularly by viral diseases (Tairo, F., *et al.*, 2005; valverde, R.A., *et al.*, 2007; Karyeija, R.F., *et al.*, 1998). Since Sweetpotato is cultivated from shoot cuttings grown in the field from previous season, percentage of virus infection increases, as consequence, total yield is dramatically reduced. The most common of these viral diseases known under the name Sweetpotato Virus Diseases (SPVD), caused by simultaneous infection with Sweetpotato Feathery Mottle Virus (SPFMV) and Sweetpotato Chlorotic Stunt Virus (SPCSV), (Winter, S., *et al.*, 1992; Gibson, R.W *et al.*, 1998).SPFMV (SPFMV; family potyviridae, genus potyvirus), is found most commonly in sweetpotatos in different parts of the world (Loebenstein G., *et al.*, 2003; Tairo, F., *et al.*, 2005). The potyvirus enhances accumulation of the unrelated virus (Karyeija, R.F., *et al.*, 2000; Latham, J.R. and A.K. Wilson, 2008). SPFMV, alone does not cause major yield losses, but co-infection and the consequent synergistic interaction of these two viruses cause the development of the sever SPVD, (Schaefer, G.A and E.R. Terry, 1976), which results in the reduction of sweetpotato yield for up to 90% while the infection of each virus alone results in reduction of up to 10% (Ishac, J., *et al.*, 2000; Carey, E.E *et al.*,1999; Gutierrez, D.L *et al.*, 2003; Njeru, R.W *et al.*, 2004; Mukasa, S.B *et al.*, 2006). Owing to the great yield losses caused

Corresponding Author: Mervat, M.M.EL FAR., Agricultural Genetic Engineering Research Institute (AGERI),
Agricultural Research Centre (ARC), 9 Gamaa St. 12619, Giza, Egypt
Email: ahmedmervat@yahoo.com

by virus diseases, their absence from propagation materials is essential for sustainable plant production. This is particularly important with vegetatively propagated crops in which infected planting materials transmit the pathogen to the following generation. Shoot tip, meristem tip and heat therapy have been widely used to produce virus-free clones of crop plants which are propagated vegetatively (Faccioli, G., and F. Marani, 1998). Sweetpotato was one of the first crops subjected to virus eradication with this technique (Alconero, R., *et al.*, 1975; Frison, E.A. and S.Y. Ng, 1981; Green, S.K., and C.Y. Lo, 1989). The previous studies (Abo El-Abbas, F., *et al.*, 1998) demonstrates the presence of SPVD in Egypt and its effect on sweetpotato production, and emphasized the need to control it by developing a system to produce virus-free plant material.

The aim of the present study is to improve the agronomic quality of deteriorated local cultivar, Abees using an efficient method for virus elimination by combining thermotherapy technique with meristem tip culture.

MATERIALS AND METHODS

Plant Material:

Sixty five sweetpotato Abees cultivar apical cuttings in a length of 15 cm were collected from several locations in Kafr El Shikh governorate, and kept under insect proof greenhouse.

Dot-ELISA:

Cuttings were tested for the presence of SPFMV using dot-ELISA following the method described by (Gutierrez, D.L *et al.*, 2003) and the modifications of (Ashoub, A. *et al* 2008). Samples were tested using antiserum raised against SPFMV, kindly provided by International Potato Centre (CIP), and a polyclonal alkaline phosphatase labelled goat anti rabbit as secondary antibodies (Sigma, USA).

Thermotherapy:

Plants were subjected to the thermotherapy treatment as follows: Plants were placed in a growth chamber under the conditions of 16 hours photoperiod at 28 °C and 8 hours dark period at 25 °C. Temperatures were raised 2 °C/ 2 days till final setting of 42 °C day and 39 °C night was obtained. Plants were maintained under these conditions for additional 4 weeks.

Meristem Tip Culture:

Following thermotherapy, axillary shoot tips with a long of 2-3 cm were cut, surfaced sterilized by immersing in 90% ethanol for 10 seconds followed by 2% sodium hypochlorite solution, 0.05% tween-20 for 5 minutes and rinsing for three times, 2 minutes each, with a sterile distilled water. A part of the apex of approximately 0.25 mm long with two leaf primordia was excised for culture on a 5-cm Petri dish containing 15 ml of MS salts medium (Murashige, T., and F. Skoog, 1962) supplemented with 3 % sucrose. The cultures were grown at 28 ± 2 °C under a 16-h photoperiod of 300 foot candle.

Virus Indexing:

Elongated shoots were tested for success of eradication of SPFMV using reverse transcriptase polymerase chain reaction, RT-PCR, as follows:

Total RNA was extracted from 50-100 mg sweetpotato leaves using SV total RNA extraction kit (Promega, USA) following the manufacturer recommendations. Reverse transcription and PCR reactions were carried out in according to the described method (Ashoub, A. *et al.*, 2008).

SPFMV-free sweetpotato plants were massive *in vitro* propagated via nodal cuttings using the media and incubation conditions described above. To assure the absence of SPFMV from the micropropagated plant materials, leaf samples were collected every 2 months and tested by dot-ELISA as described above.

RESULTS AND DISCUSSIONS

SPVD which caused by SPFMV and SPCSV is one of the most important complexes, effort to develop virus-free cultivar to one of this viruses seemed sufficient to control SPVD.

Incidence of Sweetpotato Feathery Mottle Virus (SPFMV):

Serological detection methods such as most types of ELISA lack the sensitivity required for the detection of SPFMV in sweetpotato tissue. (Esbenshade, P.R. and J.W. Moyer, 1982). However dot-ELISA is more sensitive than microplate ELISA (Abad, J.A and J.W. Moyer, 1992; Gibbs, K.S and A.C. Padovan, 1993; Fuentes, S. and L.F. Salazar, 1992).

Dot-ELISA carried out on sweetpotato plants collected from Egyptian fields showed 100% infection (fig.1), in all, 65 collected sweetpotato samples. In Egypt, Isahck, J.A. *et al.*, 2003; and F. Abo El-Abbas *et al.*, 1998) found that SPFMV is predominant in all planting material. Sweetpotato is propagated vegetatively using cuttings either produced as sprouts from tuberous roots or obtained from the foliage of mature crops (CIP, 2001), if when infected tuberous roots from a previous crop sprout in the same field or in areas adjacent to new crop, they serve as a source of virus inoculum (Karyeija, R.F., *et al.*, 1998). The majority of the farmers (96%) obtain planting material from their stocks or borrow vines from neighbours. Due to lack of knowledge, most farmers (70%) select planting vines from healthy looking vigorously growing plants (Ateka, M.E. *et al.*, 2004). However they did not recognize virus-like symptoms as a disease but associated viral symptoms with other factors such as drought, insect pests, and cultivar-specific behaviour. Lack of SPFMV symptoms as mono infection on sweetpotato plants prevents SPFMV-free plants of being selected as sources of cuttings (Clark, C.A. and J.W. Moyer, 1988; Brunt, A.A., *et al.*, 1996; Gibson, R.W. *et al.* 1997). This is probably explaining the wide distribution of the virus diseases. Absence of visual symptoms in sweetpotato is an inadequate test for SPFMV, since the virus cause slight or no symptoms when infected alone.




Fig. 1: Dot-ELISA of sweetpotato sample collected from Egyptian fields. Healthy sample is indicated in a box at the top left handsite.

Effect of Heattherapy and Meristem Culture on Survival and Regrowth:

Nearly, all (100%) of the heattherapy treated plants survived. The gradual regime used for raising the temperature allowed plants to survive. However the regrowth ability of the meristem tip decreased, only 96% of the meristem tips survived and formed shoots, when meristem tips with two leaf primordia with the size of 0.25 mm were used as plant material (fig.2-A). These results are in agreement with the results of (Faccioli, G. and F. Marani, 1998; Fuglie, K.O. *et al.*, 1999). They reported that meristem tips of 0.2 and 0.3 mm with two leaf primordia are commonly used for virus elimination in various plant species, and the survival and growth are greatly improved if the tissue taken includes slightly expanded leaf primordia.

Effect of Heattherapy Treatment and Meristem Tip Culture on Virus Elimination:

Out of sixty five SPFMV infected plants, sixty two SPFMV virus-free plants become established. (fig. 2-B). This study showed that SPFMV can be very efficiently eliminated from sweetpotato infected plants using the traditional heattherapy in a combination with meristem tip culture.(Cooper, V.C. and D.J.A. Walkey, 1978; Walkey, D.J.A. and V.C. Cooper, 1975; Hu, C.U. and P.J. Wang, 1983; Mink, J.I. *et al.*,1998;). While elimination of viruses by meristem culture is size dependent, it was recommended that the smaller the tip that is taken, the better chance for virus exclusion (data not shown), it comes with the findings of (Michael, J.K., 1996), because in the meristematic zone the rate of plant growth is increasing comparing with the rate of virus multiplication. Meristem culture technique has regenerates over 35 genera (Quak, F., 1987). In contrast (Lankes, C.1995; Karesove, R., *et al.*, 2002; Theiler-Hedtrich, R. and G. Baumann,1989) reported that meristem tip culture alone obtained a few virus-free plants. However thermotherapy followed by meristem culture resulted in SPFMV-free plants. This finding raised the question of the mechanism by which thermotherapy enhanced eradication of SPFMV. Early studies indicated that thermotherapy inhibits viral replication while virus




Fig. 2: Production of virus-free sweetpotato plants using heattherapy and meristem tip culture. (A) Apical meristem tip (0.25mm with two leaf primordia) used for virus eradication. (B) Regrowth of meristem tip after 4 weeks of culture on MS medium supplemented with 3% sucrose. (C) An established healthy plant of sweetpotato Abees cultivar obtained from SPFMV-affected plant using heattherapy and meristem tip culture.

degradation continues, which results in subsequent elimination of the virus from meristem tips (Cooper, V.C. and D.J.A. Walkey, 1978; Kassanis, B., 1975), later (Szittyá, G., *et al.*, 2003; Qu, F., *et al.*, 2005; Chellappan, P., *et al.*, 2005) found that, the efficiency of virus induced RNA silencing found to be significantly enhanced at the high temperature. Indeed, since, the first studies that aimed to produce virus-free plants from infected individuals, it has been known that the efficiency of virus eradication in a given host species differs depending on the virus and the host genotype. With some plants, thermotherapy followed by meristem tip culture did not result in any virus free plantlet. Recently, a novel approach based on cryotherapy was found to solve this problem, (Wang, Q.C. and J.P.T. Valkonen, 2008).

Indexing of Plants for SPFMV:

After heattherapy and meristem tip culture, the *in vitro* sixty two rooted plantlets two months old (fig. 2-C) subjected to RT-PCR analysis using specifically designed primers based on the conserved regions present in all viral strains obtained from the NCBI database. All detected samples were found to be SPFMV-free, a single band corresponding to the expected size of the amplified product of ~300 nucleotides. Figure (3-A) represents an example of the RT-PCR results obtained. Meanwhile, figure (3-B) showing an equal quality of RNA extracted from plants indicating that the absence of amplification in the treated samples is due to the absence of SPFMV and not to the quality of extracted RNA. Indexing method based on RT-PCR, although it is quite laborious and expensive (Fuglie, K.O. *et al.*, 1999; Valverde, R.A., *et al.*, 2007). However, it is sensitive.




Fig. 3: (A) RT-PCR of sweetpotato samples after thermotherapy and meristem tip culture. Positive control (I), one kb plus molecular weight marker, invitrogen, USA (M) and reformed sweetpotato plants (1-6). (B) Aliquot of 2µg total RNA extracted from positive control and sweetpotato samples. Arrows indicate the 28s and 18s ribosomal RNA.

Dot-ELISA was routinely used to evaluate tissue culture materials over 2 years to confirm that the absence of SPFMV from sweetpotato plants is due to the freeing of plants from infection and not that the virus is under detection level.

Conclusion:

In the present study, we report the successful elimination of SPFMV from sweetpotato the Egyptian Abees cultivar by heattherapy and meristem tip culture of infected plants.

ACKNOWLEDGMENTS

We thank the International Potato Centre (CIP), Lima, Peru, for providing the antibodies for sweetpotato virus detection. Support and partial funding was provided by the Midwest Universities Consortium for International Activities (MUCIA)/USAID project No B-24

REFERENCES

- Abad, J.A., J.W. Moyer, 1992. Detection and distribution of sweet potato feathery mottle virus in sweet potato by *in vitro*-transcribed RNA probes (riboprobes), membrane immunobinding assay, and direct blotting. *Phytopathology*, 82: 300-305.
- Abo El-Abbas, F., J. Ishak and R. El-Bedewy, 1998. Control of sweetpotato viruses. Annual Research Progress Report (0803) for Egypt. CIP, Lima, Peru, pp: 9.
- Alconero, R., A.G. Santiago, F. Morales, K.O. Fuglie, L. Zhang, L. Salazar and T. Walker, 1999. Economic Impact of virus-free sweetpotato planting material in shandong province, China, CIP program report 1997-1998.
- Alconero, R., A.G. Santiago, F. Morales, F. Rodriguez, 1975. Meristem tip culture and virus indexing of sweet potatoes. *Phytopathology*, 65: 769-773.
- Ashoub, A., Mervat, M.M.EL FAR., P. Dirk and Taymour, N. EL-DIN., 2008. Comparison of methods to detect sweetpotato feathery mottle virus (SPFMV) in sweetpotatoes.
- Ateka, M.E., R.W. Njeru, A.G. Kibaru, J.W. Kimenju, E. Barg, R.W. Gibson and H.J. Vetten, 2004. Identification and distribution of viruses infecting sweetpotato in Kenya. *Ann. Appl. Biol.*, 144: 371-379.
- Austin, D.F., 1987. The taxonomy, evolution and genetic diversity of sweetpotato and related absence of viruses in most symptomless fieldgrown sweetpotato in Uganda. *Ann. Appl. Biol.*, 130: 481-490.
- Brunt, A.A., K. Crabtree, M.J. Dallwitz, A.J. Gibbs and L. Watson, 1996. *Viruses of Plants. Description and Lists from the VIDE Database*. C.A.B. International, Wallingford, Eng.
- Carey, E.E., V. Hagenimana, M.A. Oyunga, L. K'osambo, N.E.J.M. Smit, C. Ocitti p'Obwoya, G. Turyamureeba and J. Low, 1999. Using orange-fleshed sweetpotato varieties to combat vitamin A deficiency and enhance market opportunities for smallholder farmers in sub-Saharan Africa. pp. 157-168. In: M.O. Akoroda and J.M. Teri, eds. *Food security and crop diversification in SADC countries: the role of cassava and sweetpotato. Proceedings of the scientific workshop of the Southern African Root Crops Research Network (SARRNET)*, Lusaka, Zambia, 17 - 21 August 1998. IITA, Idadan, Nigeria.
- Chellappan, P., R. Vanitharani, F. Ogbe and C.M. Fauquet, 2005. Effect of temperature on geminivirus-induced RNA silencing in plants. *Plant Physiol.*, 138: 1828-1841.
- CIP (International Potato Center), 2001. *Techniques in Plant Virology in CIP. Training Manual Version: 10 January 2001*. L.F. Salazar and U. Jayasinghe, eds. CIP, Lima, Peru.
- Clark, C.A., and J.W. Moyer, 1988. *Compendium of Sweet Potato Diseases*. American Phytopathological Society Press, St. Paul, MN.
- Cooper, V.C. and D.J.A. Walkey, 1978. Thermal inactivation of cherry leaf roll virus in tissue cultures of *Nicotiana rustica* raised from seeds and meristem tips. *Ann. Appl. Biol.*, 88: 273-278.
- Eshenshade, P.R., and J.W. Moyer, 1982. Indexing system for sweetpotato feathery mottle virus in sweetpotato using enzyme-linked immunosorbent assay. *Plant Dis.*, 66: 911-913.
- Faccioli, G., F. Marani, 1998. Virus elimination by meristem tip culture and tip micrografting. In: Hadidi, A., Khetarpal, R.K., Koganzawa, H. (Eds.), *Plant Virus Disease Control*. APS Press, St. Paul, MN, USA, pp. 346-380.
- FAOSTAT, 2006. FAO database. [<http://faostat.fao.org/site/567/default.aspx>].
- Frison, E.A., S.Y. Ng, 1981. Elimination of sweet potato virus disease agents by meristem culture. *Tropical Pest Management* 27, 452-454.

- Fuentes, S. and L.F. Salazar, 1992. Identification of a new sweetpotato virus. *Fitopatologia* 27:50 (Abstr., in Spanish).
- Fuglie, K.O., L. Zhang, L. Salazar and T. Walker, 1999. Economic Impact of virus-free sweetpotato planting material in shandong provience, China, CIP program report, 1997-1998.
- Gibbs, K.S. and A.C. Padovan, 1993. Detection of sweetpotato feathery mottle virus in sweetpotato grown in Northern Australia using an effiecient and simple assay. *Intern. J. Pest Manage*, 39: 223-228.
- Gibson, R.W., R.O.M. Mwanga, S. Kasule, I. Mpembe and E.E. Carey, 1997. Apparent.
- Gibson, R.W., I. Mpembe, T. Alicai, E.E. Carey, R.O.M. Mwanga, S.E. Seal, H.J. Vetten, 1998. Symptoms, aetiology and serological analysis of sweet potato virus disease in Uganda. *Plant Pathology*, 47: 95-102.
- Green, S.K., C.Y. Lo, 1989. Elimination of sweet potato yellow dwarf virus (SPYDV) by meristem tip culture and by heat treatment. *Journal of Plant Disease and Protection*, 96: 464-469.
- Gutiérrez, D.L., S. Fuentes, L.F. Salazar, 2003. Sweetpotato virus disease (SPVD): distribution, incidence, and effect on sweetpotato yield in Peru. *Plant Disease*, 87: 297-302.
- Hu, C.Y., P.J. Wang, 1983. In: Evans, D.A., Sharp, W.R., Ammirato P.V., Yamada, Y. (Eds.), *Handbook of Plant Cell Culture*. Macmillan, New York, pp: 177-227.
- Ishac, J., H. Bekheit, L. Ibrahim, R. El-Bedewy and F. Abo El-Abbas, 2000. Occurrence of sweetpotato virus disease in Egypt: Detection and epidemiological aspects. Fifth Triennial Congress of the African Potato Association, pp: 35-35 (Abstr.).
- Ishak, J.A., J.F. Kreuze, A. Johansson, S.B. Mukasa, F. Tairo, F.M. Abo El-Abbas and J.P.T. Valkonen, 2003. Some molecular characteristics of three viruses in sweet potato virus disease-affected sweet potato plants in Egypt. *Arch. Virol.*, 148: 2449-2460.
- Karesova, R., M. Janeckova and F. Paprstein, 2002 Elimination of Raspberry bushy dwarf virus from raspberry cv. "Gatineau". *Acta Hort.*, 585: 359-362.
- Karyeija, R.F., R.W. Gibson, J.P.T. Valkonen, 1998. The significance of sweet potato feathery mottle virus in subsistence sweet potato production in Africa. *Plant Disease*, 82: 4-15.
- Karyeija, R.F., Kreuze, J.F., Gibson, R.W., Valkonen, J.P.T., 2000. Synergistic interactions of a potyvirus and a phloem-limited crinivirus in sweet potato plants. *Virology* 269, 26-36.
- Kassanis, B., 1975. Effect of changing temperature on plant virus diseases. *Adv. Virus Res.*, 4: 221-241.
- Lankes, C., 1995. Elimination of raspberry bushy dwarf virus. *Act. Hort.*, 385: 70-75.
- Latham, J.R., A.K. Wilson, 2008. Transcomplementation and synergism in plants: implications for viral transgenes? *Molecular Plant Pathology*, 9: 85-103.
- Loebenstein, G., S. Fuentes, J. Cohen, L.F. Salazar, 2003. Sweet potato. In: Loebenstein, G., Thottappilly, G. (Eds.), *Virus and Virus-Like Diseases of Major Crops in Developing Countries*. Kluwer Academic Publishers, Dordrecht, The Netherlands, pp: 223-248.
- Michael, J., Klopmeier. 1996. Indexing for plant pathogens. In: *Plant tissue culture concepts and laboratory exercises* (CRC press LLC), pp: 341-346.
- Mink, G.I., R. Wample and W.E. Howell, 1998. Heat treatment of perennial plants to eliminate phytoplasmas, viruses, and viroids while maintaining plant survival. In: *Plant Virus Disease Control* (Hadid, A., Khetarpal, R.K. and Koganezawa, H., eds), pp: 332-345. St. Paul, MN: APS Press, The American Phytopathological Society.
- Mukasa, S.B., P.R. Rubaihayo, J.P.T. Valkonen, 2006. Interactions between a crinivirus, an impomovirus and a potyvirus in coinfection sweetpotato plants. *Plant Pathology*, 53: 458-467.
- Murashige, T., F. Skoog, 1962. A revised medium for rapid growth and bioassays with tobacco cell cultures. *Physiologia Plantarum*, 15: 473-497.
- Njeru, R.W., M.W.K. Mburu, E. Cheramgon, R.W. Gibson, E. Obudho, D. Yobera, 2004. Studies on the physiological effects of viruses on sweet potato yield in Kenya. *Annals of Applied Biology*, 145: 71-76.
- Nutrition Institute, 1995. National survey for assessment of vit.A status in Egypt in collaboration with UNICEF.
- Qu, F., X.H. Ye, G.C. Hou, S. Sato, T.E. Clemente and T.J. Morris, 2005. RDR6 has a broad-spectrum but temperature-dependent antiviral defense role in *Nicotiana benthamiana*. *J. Virol.*, 79: 15209-15217.
- Quak, F., 1987. Therapy of individual plants. In: *Viruses of Potatoes and seed-potato production* (De Bokx, J.A. and van der Want, J.P.H., eds), pp: 151-161. Wageningen: Pudoc.
- Schaefer, G.A., E.R. Terry, 1976. Insect transmission of sweet potato disease agents in Nigeria. *Phytopathology*, 66: 642-645.

Szittyá, G., D. Silhavy, A. Molnar, Z. Havelda, A. Lovas, L. Lakatos, Z. Banfalvi, and J. Burgyan, 2003. Low temperature inhibits RNA silencing-mediated defence by the control of siRNA generation. *EMBO J.*, 22: 633-640.

Tairo, F., S.B. Mukasa, R.A. Jones, A. Kullaya, P.R. Rubaihayo, J.P.T. Valkonen, 2005. Unravelling the genetic diversity of the three main viruses involved in sweetpotato virus disease (SPVD), and its practical implications. *Molecular Plant Pathology*, 6: 199-211.

Theiler-Hedtrich, R. and G. Baumann, 1989. Elimination of apple mosaic virus and raspberry bushy dwarf virus from infected red raspberry (*Rubus idaeus* L.) by tissue culture. *J. phytopath.*, 127: 193-199.

Valverde, R.A., C.A. Clark, J.P.T. Valkonen, 2007. Viruses and virus disease complexes of sweetpotato. *Plant Viruses* 1, 116-126 (1749-6209). Wang, Q.C., 1996. Factors affecting the survival and growth of in vitro grafted apple. *Journal of Fruit Science*, 13: 79-83 (in Chinese with English abstract).

Walkey, D.G.A. and V.C. Cooper, 1975. Effect of temperature on virus eradication and growth of infected tissue culture. *Ann. Appl. Biol.*, 80: 185-190.

Wang, Q.C., W.J. Cuellar, M.L. Rajamäki, Y. Hiraka, J.P.T. Valkonen, 2008. Combined thermotherapy and cryotherapy for efficient virus eradication: relation of virus distribution, subcellular changes, cell survival and viral RNA degradation in shoot tips. *Molecular Plant Pathology*, 9: 237-250.

Winter, S., A. Purac, F. Leggett, E.A. Frison, H.W. Rossel and R.I. Hamilton, 1992. Partial characterization and molecular cloning of a closterovirus from sweetpotato infected with the sweetpotato virus disease complex from Nigeria. *Phytopathology*, 82: 869-875.

Woolfe, J.A., 1992. Sweetpotato, an Untapped Food Resource. Cambridge University Press, New York. World Health Organization (WHO), 2003. Nutrition report, 16-52-28 (updated 2003)

Zhang, D., M. Chislain, Z. Huaman, A. Golmirzaie and R.J. Hijmans, 1998. RAPD variation in sweetpotato (*Ipomoea batatas*(L.)Lam.) Cultivars from South America and Papua New Guinea. *Genetic Resources and crop evolution*, 45: 271-277.